

ECO-T[®]

The fungus *Trichoderma harzianum* helps control root diseases and enhances the growth of a wide variety of vegetables, crops, ornamentals and Eucalyptus species.

Warnings

- Keep out of reach of children, uninformed persons and animals.
- Store in a refrigerator away from chemical pesticides.
- May cause slight eye irritation.
- May cause allergic reactions in sensitive persons.

Precautions

- **Remember you are working with a living organism.**
- Do not use after the expiry date.
- Do not mix in containers that have previously been used for toxic substances. Wash the containers well with soap and water before use.
- Use within 6 hours of mixing.
- Do not leave the powder or mixture standing in the sun.

- Shake or stir the fungus mixture frequently during use to avoid settling out of the product.
- Do not mix the fungus with any agrochemicals without first consulting your sales representative.
- Avoid inhaling the dust.
- Wash hands with soap and water before eating, drinking or smoking.
- Do not use under extreme conditions i.e. above 35°C or at soil/water pH less than 4 or higher than 7.

Disclaimer

Although this inoculant has been tested under a variety of conditions, the supplier does not warrant that it will be efficacious under all conditions. Because the active agent is a living organism, its action may be affected by storage, handling, climatic or soil conditions.

This naturally-occurring organism should not cause damage to crops, vegetation, or the environment, nor harm to man and animals. However, in the unlikely event of this happening, the supplier does not accept responsibility for any damage that may occur.

Eco-T[®]

Die swam *Trichoderma harzianum* help om wortelsiektes te beheer en verbeter die groei van 'n wye verskeidenheid groente, gewasse, ornamentele plante en Eucalyptus soorte.

Waarskuwings

- Hou buite bereik van kinders, oningeligte persone en diere.
- Stoor in 'n yskas (moet nie vries nie) weg van chemiese middels.
- Mag ligte oog irritasies veroorsaak.
- Mag 'n allergiese reaksie in sensitiewe persone veroorsaak.

Voorsorgmaatreëls

- **Onthou u werk met 'n lewende organisme.**
- Moet nie na die vervaldatum gebruik nie.
- Moet nie meng in houers wat voorheen vir toksiese middels gebruik was nie. Was die houers goed met seep en water voor en na gebruik.
- Gebruik binne 6 uur na vermenging.
- Moet nie die poeier of mengsel in die son laat staan nie.

- Skud of roer die mengsel dikwels gedurende gebruik.
- Moet nie met enige ander chemiese landboumiddels meng sonder om u verkoopsteenwoordiger te raadpleeg nie.
- Voorkom inaseming van die stof.
- Was hande met seep en water voordat u eet, drink of rook.
- Moet nie gebruik onder uiterste toestande nie d.w.s. bo 35°C of by grond /water pH onder 4 of bo 7.

Hoewel die inokulant onder 'n wye verskeidenheid van toestande getoets is, waarborg die verskaffer nie dat die produk effektief in alle omstandighede sal wees nie. Omdat die aktiewe bestanddeel 'n lewende organisme is, word dit soms deur berging, hanteering, klimaat en grondtoestande geaffekteer.

Dié natuurlike organisme behoort nie skade aan mense, diere, gewasse, plantegroei of die omgewing te doen nie. In die onwaarskynlikheid dat dit wel gebeur, aanvaar die verskaffer nie verantwoordelikheid daarvoor nie.

Directions for use

Seed treatment: Seed dressing application rates differ for different crops. Apply to maize at a rate of 1 g/kg of seed or 25-50 g/ha, to wheat at 1-2 g/kg seed or 40-80 g/ha, or to soybeans at 1 g/kg seed or 50-75 g/ha, together with a suitable sticker (available from the supplier). For rates on other crops please consult your local agent or Plant Health Products or www.plant-health.co.za for details.

A sticker is not required where Eco-T is added during on-farm treatment with insecticides, compatible fungicides, or Rhizobium inoculant. Apply as a dry powder to slightly moist seed.

Drench: Apply as a drench to seedling trays, seedbeds, potted plants and turf at planting, and monthly thereafter. Mix at a rate of 1 g / 4 litres water. Mix the required amount of the powder to a thin paste with a small amount of water before adding the remaining water. Irrigate the fungus into the root zone after application. The contents of this bottle treats 3000 m² of seedbed or 3000 speedling trays.

Gebruiksaanwysings

Saadbehandeling: Die dosis verskil vir die verskeie gewasse. Gebruik as 'n saadbehandeling op mielies teen 1 g/kg saad of 25-50 g/ha, op koring teen 1-2 g/kg saad of 40-80 g/ha, of sojabone teen 1 g/kg saad of 50-75 g/ha. Voeg kleefstof by om die swam op die saad vas te plak (beskikbaar van die verskaffer). Vir die dosis op ander gewasse raadpleeg asseblief jou naaste agent of Plant Health Products of www.plant-health.co.za.

Kleefstof is nie nodig wanneer Eco-T bygevoeg word gedurende plaasbehandelings met insektdoders, verenigbare swamdoders of Rhizobium entstof nie. Wend aan as 'n droë poeier op saad wat effens klam is.

Drenk: Wend aan op speedlingkassies, saadbeddens, gehouerde plante en turfgras by saai en maandeliks daarna. Meng teen een gram per 4 liter water. Meng eers die regte hoeveelheid poeier met 'n bietjie water voordat u die res van die water byvoeg. Besproei na toediening. Die inhoud van die houer behandel 3000m² saadbed of 3000 speedlingkassies.