



Registered and produced by: Plant Health Products (Pty) Ltd.
 Reg. no. 2003/007987/07
 P.O. Box 207, Nottingham Road, 3280
 Tel: +27 (0) 33 266 6130
 E-mail: info@plant-health.co.za
 Web: www.plant-health.co.za



L7945 Act 36 of 1947
 N-AR 1447
 FRAC CODE: NC

...working with nature

Marketed and distributed by: Madumbi Sustainable Agriculture (Pty) Ltd.
 Tel: +27 (0) 86 104 5261
 Email: support@madumbi.co.za
 Web: www.madumbi.co.za



A biofungicide for the protection of pruning wounds on grape vines against *Eutypa* and for the suppression of *Botrytis* on crops as specified.

Active ingredient: *Trichoderma atroviride*
 (min. 2 x 10⁹ spores per gram)

Nett Contents:

250 g	
Shipping	

Expiry Date.....

Batch no.....



CAUTION

For detailed application instructions, please refer to enclosed instructions

Eco-77®

L7945 Act No 36 of 1947
 N-AR 1447

A biofungicide for the protection of pruning wounds on grape vines against *Eutypa* and for the suppression of *Botrytis*

Active ingredient: *Trichoderma atroviride* (min. 2 x 10⁹ spores per gram)

Registered by: Plant Health Products (Pty) Ltd.
 Reg. no. 2003/007987/07
 P.O. Box 207, Nottingham Road, 3280
 Tel: +27 (0) 33 266 6130

IMPORTANT POINTS TO REMEMBER

- Remember you are working with a living organism.
- Store at 4 - 10°C for maximum shelf-life.
- Use within 6 hours of mixing.
- Do not leave the powder or mixture standing in the sun.

WARNINGS AND PRECAUTIONS

- Keep out of reach of children, uninformed persons and animals.
- May cause slight eye irritation.
- May cause allergic reactions in sensitive persons. Avoid inhaling the dust.
- Do not use after expiry date.
- Reseal packaging after use.
- Wash hands with soap and water before eating, drinking or smoking.

DIRECTIONS FOR USE - USE ONLY AS DIRECTED.

COMPATIBILITY

- Do not mix with any agrochemicals without first consulting your sales representative.
- Avoid the use of broad spectrum fungicides for at least three days before and after Eco-77 applications.

MIXING AND APPLICATION INSTRUCTIONS

- Mix with water at the recommended rate and apply as a full cover spray.
- Do not mix in containers that have previously been used for toxic substances.
- Wash mixing containers well with soap and water before use.
- Good spray coverage is essential.
- Shake or stir the fungus mixture frequently during use to avoid settling out of the product.
- Do not apply when temperatures are below 15°C or above 35°C.
- Avoid pH extremes (less than 3 or higher than 8).

REGISTERED USAGES:

Pest / Crop	Dosage	Remarks
Eutypa Grape vine pruning wounds	Apply at a rate of 1 g per 2 litres water.	Apply as a directed spray onto each pruning wound immediately after pruning.
Botrytis Tunnel or greenhouse tomato and cucumber	Apply at a rate of 1 g per 2 litres water.	Apply as a full cover spray when disease is first noticed. Repeat as needed.
Botrytis Table grapes	Apply at a rate of 1 g per 2 litres water.	1. A minimum of 250 g/ha is recommended. 2. Apply first application at full flowering, repeat at pea-berry, veraison and 7 days pre-harvest.

DISCLAIMER

Although this product has been tested under a variety of conditions, the supplier does not warrant that it will be efficacious under all conditions. Because the active agent is a living organism, its action may be affected by storage, handling, climatic or soil conditions. The supplier is also not responsible for poor performance of the product due to failure of the user to follow the label instructions.

This naturally-occurring organism should not cause damage to crops, vegetation, or the environment, nor harm to man and animals. However, in the unlikely event of this happening, the supplier does not accept responsibility for any damage that may occur.