



**Eco-T Ezi-Flo<sup>®</sup>**



Registered and produced by: Plant Health Products (Pty) Ltd.  
Reg. no. 2003/007987/07  
P.O. Box 207, Nottingham Road, 3280  
Tel: +27 (0) 33 266 6130  
E-mail: info@plant-health.co.za  
Web: www.plant-health.co.za

L9276 Act 36 of 1947  
N-AR 1618

FRAC CODE: NC

A lubricant to be added to mechanical planters when planting treated or untreated seed, with the added benefit of Eco-T, a fungal inoculant for the control of crop root diseases and for enhanced plant growth.

**Active ingredient:** *Trichoderma asperellum*  
(min. 2 x 10<sup>9</sup> spores per gram)

...working with nature

Marketed and distributed by: Madumbi Sustainable Agriculture (Pty) Ltd.



Tel: +27 (0) 86 104 5261  
Email: support@madumbi.co.za  
Web: www.madumbi.co.za

**Nett Contents:**

|          |  |
|----------|--|
| 1 kg     |  |
| Shipping |  |

Expiry Date.....



Batch no.....



**CAUTION**

*For detailed application instructions, please refer to enclosed instructions*



**Eco-T Ezi-Flo<sup>®</sup>**

L9276 Act No 36 of 1947  
N-AR 1618

A lubricant to be added to mechanical planters when planting treated or untreated seed, with the added benefit of Eco-T, a fungal inoculant for the control of crop root diseases and for enhanced plant growth.

**Active ingredient:** *Trichoderma asperellum* (min. 2 x 10<sup>9</sup> spores per gram)

Registered by: Plant Health Products (Pty) Ltd.  
Reg. no. 2003/007987/07  
P.O. Box 207, Nottingham Road, 3280  
Tel: +27 (0) 33 266 6130

**IMPORTANT POINTS TO REMEMBER**

- Remember you are working with a living organism.
- Store at 4 - 10°C for maximum shelf-life.
- Do not leave powder or mixture standing in the sun.
- Use within 6 hours of mixing.

**COMPATIBILITY**

- Do not mix the fungus with any agrochemicals without first consulting your sales representative.

**WARNINGS AND PRECAUTIONS**

- Keep out of reach of children, uninformed persons and animals.
- May cause slight eye irritation.
- May cause allergic reactions in sensitive persons.
- Do not use after the expiry date.
- Do not mix in containers that have previously been used for toxic substances.
- Wash mixing containers well with soap and water before use.
- Shake or stir the fungus mixture frequently during use to avoid settling out of the product.
- Avoid inhaling the dust.
- Wash hands with soap and water before eating, drinking or smoking.
- Do not use under extreme conditions i.e. above 35°C or at soil/water pH less than 4 or higher than 7.

**DIRECTIONS FOR USE**

| CROP   | DOSAGE RATE  | REMARKS  |
|--|--|--|
| Row crops such as maize, soya, sunflower, dry bean | 1 g per kg seed<br>Min. 25 g per hectare (1 scoop) | Apply at planting. Mix directly into seed hopper of planter. Fill half the hopper with seed, add and mix in half the product. Add the balance of the seed and mix in the balance of the product. |
| Root vegetable crops such as carrots and onions    | 50 g per kg seed (2 scoops)                        |  |

**DISCLAIMER**

Although this product has been tested under a variety of conditions, the supplier does not warrant that it will be efficacious under all conditions. Because the active agent is a living organism, its action may be affected by storage, handling, climatic or soil conditions. The supplier is also not responsible for poor performance of the product due to failure of the user to follow the label instructions.

This naturally-occurring organism should not cause damage to crops, vegetation, or the environment, nor harm to man and animals. However, in the unlikely event of this happening, the supplier does not accept responsibility for any damage that may occur.